

29 September 2000 1ED220:0CO:SRE/rlk GI-5324 P2 PATENT

- 220 -

5

10

 $R_6$  is selected from H,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy, -NO<sub>2</sub>, -CF<sub>3</sub>, or -OH;

R<sub>7</sub> is selected from -(CH<sub>2</sub>)<sub>n</sub>-COOH, -(CH<sub>2</sub>)<sub>n</sub>-N-(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -(CH<sub>2</sub>)<sub>n</sub>-NH-(C<sub>1</sub>-C<sub>6</sub> alkyl), -CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, -NH-(C<sub>1</sub>-C<sub>6</sub> alkyl), -N-(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, -NO<sub>2</sub>, -CF<sub>3</sub>, or -OH;